Bioremediation of Environmental Pollutions by Glycosylation with Immobilized Marine Microalga

Kei Shimoda ¹, Hiroki Hamada ², Yushi Uchimura ³

¹ Faculty of Medicine, Oita University; ² Faculty of Science, Okayama University of Science; ³ Ehime Prefectural Fisheries Experimental Station

Summary

Biphenyl compounds such as bisphenol A and benzophenone have been listed among "chemicals suspected of having endocrine disrupting effects" by the World Wildlife Fund, the National Institute of Environmental Health Sciences in the USA and the Japanese Environment Agency. The biphenyl compounds exhibited estrogenic activity in bioassays. These compounds are widely used to manufacture polyacrylates, ether resins, phenol resins, insecticides, agricultural chemicals, pharmaceuticals, and coatings, and their residues are released as pollutants into rivers and seas. In the case of bisphenol A, worldwide production capacity was estimated about 1,100 million pounds. From the viewpoint of seawater pollution control, metabolism of biphenyl compounds by marine microalgae is of importance.

This study focused on the metabolism of biphenyl compounds by cultured and immobilized marine microalga. Biotransformation product was isolated from the cultured cells of *A. crassum*, which had been treated with benzophenone. Potentially glucosylated product was obtained, and its chemical structure was determined on the basis of their FABMS, 1 H and 13 C NMR, H-H COSY, C-H COSY, and HMBC spectra as diphenylmethyl β -D-glucoside. The glucosylation activity of immobilized *A. crassum* cells was enhanced at 2% sodium alginate concentration. Use of immobilized *A. crassum* cells improved the glycosylation activity, and the amount of glycosides was much enhanced in comparison with the case of biotransformation with cultured *A. crassum* cells.

From the viewpoint of seawater pollution control, glycosylation of biphenyl compounds by immobilized marine microalga is important and environmentally friendly.